

**UNCLASSIFIED**

---

**AD**

**401 541**

*Reproduced  
by the*

**DEFENSE DOCUMENTATION CENTER**

**FOR**

**SCIENTIFIC AND TECHNICAL INFORMATION**

**CAMERON STATION, ALEXANDRIA, VIRGINIA**



---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

633-2

USNRDL-TR-625  
4 March 1963

CATALOGED BY ASTIA  
AS AD NO. 401541

DIFFERENTIAL EFFECTS OF SPECIFIC ANTISERA ON  
THE REJECTION OF ALLOGENIC AND XENOGENIC SKIN GRAFTS BY  
SUBLETHALLY X-IRRADIATED MICE

by  
M. L. Tyan  
L. J. Cole

APR 16 1963  
ASTIA  
A

401 541

U.S. NAVAL RADIOLOGICAL  
DEFENSE LABORATORY  
SAN FRANCISCO 24, CALIFORNIA

12ND. P7463

EXPERIMENTAL PATHOLOGY BRANCH

L. J. Cole, Head

BIOLOGICAL AND MEDICAL SCIENCES DIVISION

E. L. Alpen, Head

---

ADMINISTRATIVE INFORMATION

This work was accomplished under the Bureau of Medicine and Surgery Task MR005.08-5200, Subtask 3, Technical Objective AW-6, as described in the U. S. Naval Radiological Defense Laboratory Annual Report to the Bureau of Medicine and Surgery (OPNAV Form 3910-1) of 31 December 1962, and is listed in the U.S. Naval Radiological Defense Laboratory Technical Program Summary for Fiscal Years 1963-1965 of 1 November 1962 under Program A3, Problem 2, entitled "Nuclear Warfare Aspects of Whole Body Ionizing Radiation." This study was supported through funds provided by the Bureau of Medicine and Surgery, and the Defense Atomic Support Agency under NWER Program A4c, Subtask 03.027.

---

*Eugene P. Cooper*

Eugene P. Cooper  
Scientific Director

*E. B. Roth*

E. B. Roth, CAPT USN  
Commanding Officer and Director

## ABSTRACT

Twelve to fourteen week old female  $LAF_1$  mice, non-sensitized or pre-sensitized with two consecutive BALB/c or rat skin grafts, received 670 rad whole body X radiation and orthotopic tail grafts of  $LAF_1$ , BALB/c, C3D/2 and rat skin. Certain groups of mice received specific antisera intraperitoneally, produced in response to BALB/c or rat spleen cells or to two consecutive BALB/c or rat skin grafts. The data indicate that the first-set response to an allogenic skin graft can be significantly inhibited, in sublethally irradiated mice, by specific antisera, while the first-set response to a xenogenic skin graft remains resistant to similar treatment. Specific antisera had no effect upon a pre-existing second-set response. The significance of these data is discussed.

## SUMMARY

### The Problem:

Under certain conditions, the immunological response of mice to foreign solid tissue grafts or to particular tumors can be inhibited by the administration of specific antisera produced in response to the same tissue or tumor. It is now generally accepted that at least one effector of this passively transferred "enhanced state" is circulating antibody. To date, attempts at "enhancing" the survival of normal solid tissue grafts have been confined primarily to grafts between certain strains of mice. The present communication deals with the effects of specific antisera upon allogenic (different strain of mouse) and xenogenic (different species) skin graft survival in non-sensitized and pre-sensitized sublethally irradiated mice.

### The Findings:

The findings are described in the Abstract paragraph of this report.

## INTRODUCTION

The discovery of the phenomenon of "enhancement" of tumor growth, later shown to be passively transferred with serum from the pre-treated host (1-5), and its subsequent application to the transplantation of normal allogenic solid tissues (6-9), has provided the investigator with a potentially powerful tool for the "dissection" of the mammalian immune system. It is now generally accepted that at least one effector of the passively transferred "enhanced state" is circulating antibody, whether induced in response to lyophilized or viable tissue, neoplastic or normal (4,5).

It has been shown in mice (9) that high doses of specific antisera tend to prolong the survival of allogenic skin grafts, inhibit the development or expression of "homograft sensitivity" in response to certain antigenic stimuli, but have no effect upon a pre-existing second-set response. On the basis of their observations, Brent and Medawar (9) have postulated that "antiserum probably affects the process of sensitization itself, acting by a 'central' inhibition of unknown character rather than by obstructing the afferent or efferent pathways of response".

To date, attempts to promote the survival of homografts of normal tissues have been confined to allogenic grafts between certain strains of mice or other rodents. The present communication deals with the effects of specific antisera upon allogenic (H-2 difference) and xenogenic (rat) skin graft survival in non-sensitized and pre-sensitized,

sublethally irradiated (670 rad) mice.

Evidence will be presented demonstrating that the first-set response to an allogenic skin graft can be significantly inhibited, in sublethally irradiated mice, by passively transferred specific antisera, while the first-set response to a xenogenic skin graft remains resistant to similar treatment. Further, it will be shown that specific antisera, in the manner given, have no effect upon a pre-existing second-set response. The significance of these data will be discussed.

#### MATERIALS AND METHODS

Twelve to 14 week old female (C57L x A) $F_1$ , (LAF $_1$ ) mice were used as skin graft recipients. Skin graft donors were adult female LAF $_1$  (H2 ab), male BALB/c (H2 d) and (C3H x DBA/2) $F_1$ , (C3D/2), (H2 Kd) mice, and 2-3 week old male and female Sprague-Dawley rats. The orthotopic tail skin grafting method of Bailey and Usoma was used (10). The details of grafting and the criterion of rejection (complete destruction of the engrafted tissue) have been reported previously (11). Mean survival time of the grafts and standard deviation (S.D.) are reported.

Non-sensitized LAF $_1$  mice received 670 rad whole body X radiation followed by 0.9-2.0 ml antiserum intraperitoneally, and were then grafted with LAF $_1$ , BALB/c, C3D/2 and rat skin within six hours thereafter. Other mice were pre-sensitized with two consecutive BALB/c or rat skin grafts, and one week following the rejection of the second skin graft, they received 670 rad whole body X radiation, 1.0-1.1 ml antiserum



intraperitoneally, and were grafted as described above. The radiation factors (250 KVP, 15 ma; HVL 1.5 mm Cu; 30 rad/min) and details of exposure were the same as previously reported from this Laboratory (12).

The antisera were prepared in adult A/HeJ, C3H or LAF<sub>1</sub> mice of both sexes (see tables for particulars), either by means of three intraperitoneal injections, within nine days, of an homogenate of BALB/c or rat spleen cells (1/5 or 1/20 spleen respectively), or with two consecutive BALB/c or rat skin grafts. One week following the last injection of spleen cells or the rejection of the second skin graft, the mice were sacrificed, and the antisera were prepared from blood aspirated aseptically from the inferior vena cava. It was stored at -15°C until used. In one experiment, antisera were pooled from the three strains of mice which served as donors.

A previous communication (13) postulated the presence at the height of a vigorous second-set response of a non-specific agent which disrupts capillary integrity at the graft site, whether it be allogenic, xenogenic or isogenic. Therefore, "anti-plasma" was prepared from blood aspirated into Heparin-wetted syringes (Heparin USP; 10 mg per ml), from adult A/HeJ mice which were at the height of rejection of their third consecutive BALB/c or rat skin graft. The "anti-plasma" was administered immediately to the recipient mice (1.0-1.3 ml intraperitoneally).

It had been noted previously (11, 13) that non-irradiated and

sublethally irradiated (670 rad) mice, pre-sensitized with rat skin grafts, rejected subsequent first-set allogenic grafts significantly sooner than did their appropriate controls. It was postulated that this phenomenon represented a quantitatively expanded first-set response rather than a "non-specific" or "overflow" response to rejection of the rat skin graft (it was demonstrated previously that BALB/c or C3D/2 cells and rat skin share no common transplantation antigens with respect to LAF<sub>1</sub> mice (11)). Therefore, one group of mice pre-sensitized with rat skin grafts received 670 rad whole body X radiation, and all grafts except rat skin were placed. A second group, similarly pre-sensitized and irradiated, received 1.1 ml anti-BALB/c spleen serum intraperitoneally and all grafts were placed.

All mice were housed 10 per cage. The diet was Purina Lab Chow, and water containing 1% Neomycin was given ad lib.

## RESULTS

### First-Set Response Table I

All non-sensitized, sublethally irradiated LAF<sub>1</sub> mice given anti-BALB/c sera, produced in response to dissociated spleen cells or skin grafts, rejected subsequent BALB/c and C3D/2 skin grafts significantly later than did their appropriate controls (m.s.t. 29.7-40 days versus 21.1 days). Nine of 24 mice, so treated, rejected concurrent rat skin grafts somewhat later than did the control group (m.s.t. 27.2 days versus 22.7 days). However, only four of 15 mice given anti-rat sera

TABLE I

REJECTION OF ALLOGENIC AND XENOGENIC SKIN GRAFTS BY  
SUBLETHALLY IRRADIATED (670 RAD) LAF<sub>1</sub> MICE GIVEN ANTI-BALB/c OR ANTI-RAT SERA

TYPE OF ANTISERUM	SOURCE	AMT ML	NO. MICE	ISOGRAFTS LOST	TIME FOR COMPLETE REJECTION MEAN SURVIVAL(DAYS) ± S.D.		
					BALB/c	S3D/2	RAT
None			25	2	21.0 ± 3.5	21.1 ± 3.3	22.7 ± 3.5
Anti-BALB/c spleen cells	A/HeJ	2.0	2	0	40	32.5	26.5
Anti-BALB/c spleen cells	A/HeJ	1.0	8	1	29.1 ± 1.4	35.5 ± 12.2	27.2 ± 5.0
Anti-BALB/c skin grafts	A, C3H, LAF <sub>1</sub> *	1.0	3	1	35.0 ± 1.4	34.0 ± 3.4	19.6 ± 4.4
Anti-BALB/c skin grafts	A/HeJ	1.0	10	0	29.9 ± 3.1	29.7 ± 3.1	22.0 ± 1.7
Anti-BALB/c skin grafts	A/HeJ*	1.0	1	0	30.0	30.0	30.0
Anti-rat spleen cells	A/HeJ	1.0	2	0	22.5	22.5	23.5
Anti-rat spleen cells	LAF <sub>1</sub>	0.9	4	0	24.2 ± 2.8	23.0 ± 3.4	24.0 ± 6.0
Anti-rat skin grafts	A, C3H, LAF <sub>1</sub>	1.0	5	1	18.0 ± 5.9		22.0 ± 0.0
Anti-rat skin grafts	A/HeJ	1.3	4	0	29.0 ± 8.0	25.0 ± 7.3	28.1 ± 5.6

\* Pooled antisera

\*Heparinized plasma collected at the height of rejection of the third consecutive skin graft and used immediately.

failed to reject rat skin grafts at the expected time (i.e., control:  $22.7 \pm 3.5$  days). These mice had received anti-rat "plasma" and they also manifested a delay in rejection of concurrent allogenic skin grafts. All other mice receiving anti-rat sera rejected subsequent allogenic grafts at the expected time. At no time was there evidence either of an accelerated rejection of allogenic or xenogenic skin grafts or an increase in capillary fragility at the graft sites.

#### Second Set Response Table II

The second-set response of sublethally irradiated mice, pre-sensitized with BALB/c or rat skin grafts, was unaffected by antisera in the dosage given.

Sublethally irradiated mice, pre-sensitized with rat skin grafts, rejected subsequent allogenic skin grafts significantly sooner than did the control group despite the absence of a rat skin graft. However, when anti-BALB/c serum was given to a group of mice, similarly treated but receiving all grafts, the mean survival time of the allogenic skin grafts doubled (m.s.t. 31.6, 27.2 days versus 13.7, 14.3 days), i.e., allogenic skin graft survival was similar to that seen in the non-sensitized mice given anti-BALB/c sera, or as seen in sublethally irradiated mice previously sensitized with BALB/c spleen cells (13).

#### DISCUSSION

These and other data (9) clearly indicate that antisera, produced in response to normal allogenic tissues and in a manner which would be

TABLE II  
REJECTION OF ALLOGENIC AND XENOGENIC SKIN GRAFTS BY SUBLETHALLY IRRADIATED (670 RAD) LAF<sub>1</sub> MICE  
PRE-SENSITIZED WITH BALB/c OR RAT SKIN GRAFTS AND GIVEN ANTI-BALB/c OR ANTI-RAT SERA

MEANS OF SENSITIZATION	TYPE OF ANTISERUM	AMT. ML.	NO. MICE	ISOGRAFTS LOST	TIME FOR COMPLETE REJECTION MEAN SURVIVAL(days) ± S.D.		
					BALB/c	C3D/2	RAT
BALB/c skin grafts	anti-BALB/c spleen	1.0	7	2	11.1 ± 1.8	11.5 ± 1.0	20.2 ± 1.7
Rat skin grafts	anti-rat spleen	1.0	7	2	18.8 ± 8.7*	16.0 ± 9.1*	4.3 ± 0.6
Rat skin grafts <sup>#</sup>	none		7	0	13.7 ± 3.8	14.3 ± 2.3	none
Rat skin grafts	anti-BALB/c spleen	1.1	9	0	31.6 ± 6.0	27.2 ± 3.8	5.7 ± 1.3

\* 1/7 BALB/c skin grafts rejected at 8 days.

\* 3/7 C3D/2 skin grafts rejected between 6-8 days.

<sup>#</sup> These mice were not grafted with rat skin after irradiation.

expected to induce a state of "sensitivity" in the host, are capable of significant inhibition of the first-set response to subsequent allogenic skin grafts. However, the second-set phenomenon appears to be resistant to the action of the same antisera. These observations tend to support the hypothesis of "central inhibition" as the means by which specific antisera induce the "enhanced state".

However, neither the first nor the second-set response to a xenogenic skin graft appears to be affected by the presence of relatively large amounts of specific antisera. These and other data (11, 13) argue most strongly for the existence within the "immune system" of functionally distinct cell lines or systems, i.e., xenogenic solid tissue grafts provoke a response from a "cell line" functionally and perhaps phylogenetically different from that "cell line" which would respond to an allogenic skin graft. Rat skin does possess, in addition, other antigens which stimulate cell lines capable of reacting to allogenic skin grafts. This was demonstrated when the accelerated rejection of allogenic skin grafts by sublethally irradiated mice previously sensitized with rat skin grafts was abrogated by means of a specific antiserum (anti-BALB/c). This phenomenon most likely represents a quantitatively expanded, "radioresistant" first-set response.

A previous communication (13) has suggested the uniqueness of the "sensitivity" evoked by skin grafts, allogenic or xenogenic, as opposed to that induced by other means (see also 9, 14). It was demonstrated that the "homograft sensitivity" elicited by skin grafts was markedly

more radioresistant than was that produced with dissociated cells. Further, sensitization with allogenic dissociated spleen cells resulted in a prolongation of survival of subsequent allogenic skin grafts in sublethally irradiated mice when compared to the appropriate controls, i.e., "self enhancement".

On the basis of the cited data, it is suggested that the second-set response is indeed unique. It is postulated that the second-set response to an allogenic skin graft represents the immunological response of a minimum of two cell types or systems: first, a "monitor cell" (perhaps thymic in origin) responds to the incoming antigenic stimulus, bears prime responsibility for humoral antibody production either directly or through collateral cells, and is responsive to circulating antibody levels, i.e., positive feed-back mechanism; and second, an "effector cell" is activated and "armed" by the "monitor cell". The latter cell line or system remains "armed" and accounts for the persistence of the second-set response. Similarly, the second-set response to a xenogenic skin graft may represent a two stage response contained within a primigenial cell line or system, or perhaps, be the manifestation of two separate biochemical steps within a single cell. It is realized that the above is most speculative and that even if it represented a first approximation to the existing nature of things, many ancillary steps and auxiliary cell systems (macrophages, polymorphonuclear cells, etc.) involved in the rejection of a skin graft

have been neglected for the purposes of this discussion.

However, within this hypothetical frame-work "enhancement" of graft survival would be the resultant of prior stimulation of the "monitor cell" without concomitant "arming" of the "effector cell"; that is, the stimulus would be such as to call forth the production of humoral antibodies exclusively, and the "monitor cells" being sensitive to the level of antibody would fail to respond appropriately to a second stimulus in the form of a homograft. Passive enhancement would similarly be explained. Further, the "sensitivity," evoked by dissociated cells appears to represent an expansion within the "line of monitor cells"; the accelerated rejection of subsequent skin grafts is a manifestation of this quantitative change.



#### BIBLIOGRAPHY

1. Flexner, S. and Jobling, J. W., Proc. Soc. Exp. Biol. Med. N. Y., 4: 156, 1907.
2. Casey, A. E., Cancer Res., I: 134, 1941.
3. Snell, G. D., Cloudman, A. M. and Woodworth, E., Cancer Res. 8: 429, 1948.
4. Kaliss, N. Ann. N. Y. Acad Sci. 64: 977, 1957.
5. Kaliss, N. Cancer Res. 18: 992, 1958.
6. Billingham, R. E. and Sparrow, E. M., J. Embryol. Exper. Morph. 3: 265, 1955.
7. Billingham, R. E., Brent, L. and Medawar, P. B., Transpl. Bull. 3: 84, 1956.
8. Parkes, A. S., Transpl. Bull. 5: 45, 1958.
9. Brent, L. and Medawar, P. B., Proc. Roy. Soc. B, 155: 392, 1961.
10. Bailey, D. W. and Usoma, B., Transpl. Bull. 7: 424, 1960.
11. Tyan, M. L. and Cole, L. J. In Press.
12. Cole, L. J., Fishler, M. C., Ellis, M. E. and Bond, V. P. Proc. Soc. Exper. Biol. and Med. 80: 112, 1952.
13. Tyan, M. L. and Cole, L. J. In Press.
14. Billingham, R. E., Brent, L., Brown, J. B. and Medawar, P. B., Transpl. Bull. 4: 410, 1959.

INITIAL DISTRIBUTION

Copies

NAVY

3	Chief, Bureau of Ships (Code 335)
1	Chief, Bureau of Ships (Code 320)
2	Chief, Bureau of Medicine and Surgery
1	Chief of Naval Operations (Op-07T)
1	Chief of Naval Research (Code 104)
3	Director, Naval Research Laboratory (Code 2021)
1	Office of Naval Research (Code 422)
1	Office of Naval Research (Code 441)
10	Office of Naval Research, FPO, New York
3	Naval Medical Research Institute
1	OIC, Radiation Exposure Evaluation Laboratory
1	Director, Aviation Medical Acceleration Laboratory
1	U. S. Naval Postgraduate School, Monterey
1	Commander, Naval Ordnance Laboratory, Silver Spring
1	Naval Missile Center (Code 5700)
1	U. S. Naval Hospital, San Diego
1	CO, Naval Medical Research Unit No. 2
1	CO, Naval Medical Field Research Laboratory, Camp Lejeune

ARMY

1	Chief of Research and Development (Atomic Division)
1	Chief of Research and Development (Life Science Division)
1	Deputy Chief of Staff for Military Operations (CBR)
1	Chief of Engineers (ENGMC-DE)
1	Chief of Engineers (ENG CW)
1	CG, Army Materiel Command (AMCRD-DE-NE)
1	CG, USA CBR Agency
3	CO, BW Laboratories
1	CO, Fort McClellan Alabama
1	Commandant, Chemical Corps Schools (Library)
1	CG, CBR Combat Developments Agency
1	CO, Chemical Research and Development Laboratories
1	Commander, Chemical Corps Nuclear Defense Laboratory
1	Hq., Army Environmental Hygiene Agency
1	CG., Aberdeen Proving Ground

1 CO, Army Medical Research Laboratory  
 1 Army Medical Research and Nutrition Laboratory (MEDEN-AD)  
 1 CO, Army Medical Service Combat Developments Agency  
 2 Medical Field Service School, Fort Sam Houston (Stimson Lib.)  
 1 Brooke Army Medical Center (Dept. Prev. Med.)  
 1 Director, Surgical Research Unit, Fort Sam Houston  
 1 Director, Walter Reed Army Medical Center  
 1 Hq., Army Nuclear Medicine Research Detach., Europe  
 1 CG, Combat Developments Command (CDCMR-V)  
 1 CG, Quartermaster Res. and Eng. Command  
 1 Hq., Dugway Proving Ground  
 3 The Surgeon General (MEDNE)  
 1 Office of the Surgeon General (Combat Dev.)  
 1 CG, Engineer Res. and Dev. Laboratory  
 1 Director, Office of Special Weapons Development  
 1 CG, Munitions Command  
 1 CO, Frankford Arsenal  
 1 CG, Army Missile Command

#### AIR FORCE

1 Assistant Chief of Staff, Intelligence (AFCIN-3B)  
 6 CG, Aeronautical Systems Division (ASAPRD-NS)  
 1 CO, Radiological Health Laboratory Division  
 1 Director, USAF Project RAND  
 1 Commandant, School of Aerospace Medicine, Brooks AFB  
 1 CO, School of Aviation Medicine, Gunter AFB  
 1 6571st Aeromedical Research Lab., Holloman AFB  
 1 Radiobiological Laboratory  
 1 Office of the Surgeon (SUP3.1), Strategic Air Command  
 1 Office of the Surgeon General  
 1 CG, Special Weapons Center, Kirtland AFB  
 1 Director, Air University Library, Maxwell AFB  
 2 Commander, Technical Training Wing. 3415th TTG  
 1 Hq., Second Air Force, Barksdale AFB  
 1 Commander, Electronic Systems Division (CRZT)

#### OTHER DOD ACTIVITIES

3 Chief, Defense Atomic Support Agency (Library)  
 1 Commander, FC/DASA, Sandia Base (FCDV)  
 1 Commander, FC/DASA, Sandia Base (FCTG5, Library)  
 1 Commander, FC/DASA, Sandia Base (FCWT)  
 2 Office of Civil Defense, Washington  
 2 Civil Defense Unit, Army Library  
 1 Armed Forces Institute of Pathology  
 20 Armed Services Technical Information Agency  
 1 Director, Armed Forces Radiobiology Research Institute

# AEC ACTIVITIES AND OTHERS

1 Research Analysis Corporation  
1 Life Science Officer, AEC, Washington  
1 Director, Division of Biology and Medicine  
1 NASA, Ames Research Center, Moffett Field  
1 Naval Attache, Stockholm (for Commodore Troell)  
1 Aerojet General, Azusa  
5 Argonne Cancer Research Hospital  
10 Argonne National Laboratory  
2 Atomic Bomb Casualty Commission  
1 AEC Scientific Representative, France  
1 AEC Scientific Representative, Japan  
3 Atomic Energy Commission, Washington  
2 Atomic Energy of Canada, Limited  
3 Atomics International  
2 Battelle Memorial Institute  
1 Borden Chemical Company  
3 Brookhaven National Laboratory  
1 Chicago Patent Group  
1 Colorado State University  
1 Columbia University (Rossi)  
1 Committee on the Effects of Atomic Radiation  
3 Defence Research Member  
2 duPont Company, Aiken  
1 duPont Company, Wilmington  
1 Edgerton, Germeshausen and Grier, Inc., Goleta  
1 Edgerton, Germeshausen and Grier, Inc., Las Vegas  
2 General Dynamics, Fort Worth  
2 General Electric Company, Cincinnati  
8 General Electric Company, Richland  
1 General Electric Company, St. Petersburg  
1 General Scientific Corporation  
1 Hughes Aircraft Company, Culver City  
1 Iowa State University  
1 Journal of Nuclear Medicine  
1 Knolls Atomic Power Laboratory  
2 Los Alamos Scientific Laboratory (Library)  
1 Lovelace Foundation  
1 Martin-Marietta Corporation  
1 Massachusetts Institute of Technology  
1 Mound Laboratory  
1 National Academy of Sciences  
2 NASA, Scientific and Technical Information Facility  
1 National Bureau of Standards (Taylor)  
1 National Cancer Institute  
1 National Lead Company of Ohio  
1 National Library of Medicine  
1 New Jersey State Department of Health

1 New York Operations Office  
 1 New York University (Eisenbud)  
 1 Office of Assistant General Counsel for Patents  
 2 Phillips Petroleum Company  
 4 Pratt and Whitney Aircraft Division  
 2 Public Health Service, Washington  
 1 Public Health Service, Las Vegas  
 1 Public Health Service, Montgomery  
 1 Sandia Corporation, Albuquerque  
 1 Union Carbide Nuclear Company (ORGDP)  
 5 Union Carbide Nuclear Company (ORNL)  
 1 Union Carbide Nuclear Company (Paducah Plant)  
 1 United Nuclear Corporation (NDA)  
 1 U. S. Geological Survey, Denver  
 1 U. S. Geological Survey, Menlo Park  
 1 U. S. Geological Survey, Naval Gun Factory  
 1 U. S. Geological Survey, Washington  
 1 U. S. Weather Bureau, Washington  
 1 University of California, Davis  
 3 University of California Lawrence Radiation Lab., Berkeley  
 2 University of California Lawrence Radiation Lab., Livermore  
 1 University of California, Los Angeles  
 1 University of California, San Francisco  
 1 University of Chicago Radiation Laboratory  
 1 University of Hawaii  
 1 University of Puerto Rico  
 1 University of Rochester (Atomic Energy Project)  
 1 University of Tennessee (UTA)  
 1 University of Utah  
 1 University of Washington (Donaldson)  
 1 Wayne State University  
 1 Westinghouse Electric Corporation (Rahilly)  
 1 Westinghouse Electric Corporation (NASA)  
 1 Western Reserve University (Friedell)  
 25 Technical Information Extension, Oak Ridge

USNRDL

41 USNRDL, Technical Information Division

DISTRIBUTION DATE: 27 March 1963

<p>Naval Radiological Defense Laboratory USNRDL-TR-625</p> <p>DIFFERENTIAL EFFECTS OF SPECIFIC ANTISERA ON THE REJECTION OF ALLOGENIC AND XENOGENIC SKIN GRAFTS BY SUBLETHALLY X-IRRADIATED MICE by M. L. Tyan and L. J. Cole 4 March 1963 19 p. tables 14 refs.</p> <p>UNCLASSIFIED</p> <p>Twelve to fourteen week old female LAF<sub>1</sub> mice, non-sensitized or pre-sensitized with two consecutive BALB/c or rat skin grafts,</p> <p>(over)</p>	<p>Naval Radiological Defense Laboratory USNRDL-TR-625</p> <p>DIFFERENTIAL EFFECTS OF SPECIFIC ANTISERA ON THE REJECTION OF ALLOGENIC AND XENOGENIC SKIN GRAFTS BY SUBLETHALLY X-IRRADIATED MICE by M. L. Tyan and L. J. Cole 4 March 1963 19 p. tables 14 refs.</p> <p>UNCLASSIFIED</p> <p>Twelve to fourteen week old female LAF<sub>1</sub> mice, non-sensitized or pre-sensitized with two consecutive BALB/c or rat skin grafts,</p> <p>(over)</p>	<p>1. Immune serums. 2. Skin. 3. Transplantation. 4. Radiation tolerance. 5. Mice. 6. Rats.</p> <p>I. Tyan, M. L. II. Cole, L. J. III. Title. IV. MR005.08-5200</p> <p>UNCLASSIFIED</p> <p>received 670 rad whole body X radiation and orthotopic tail grafts of LAF<sub>1</sub>, BALB/c, C3D/2 and rat skin. Certain groups of mice received specific antisera intraperitoneally, produced in response to BALB/c or rat spleen cells or to two consecutive BALB/c or rat skin grafts. The data indicate that the first-set response to an allogenic skin graft can be significantly inhibited, in sublethally irradiated mice, by specific antisera, while the first-set response to a xenogenic skin graft remains resistant to similar treatment. Specific antisera had no effect upon a pre-existing second-set response. The significance of these data is discussed.</p> <p>UNCLASSIFIED</p>	<p>1. Immune serums. 2. Skin. 3. Transplantation. 4. Radiation tolerance. 5. Mice. 6. Rats.</p> <p>I. Tyan, M. L. II. Cole, L. J. III. Title. IV. MR005.08-5200</p> <p>UNCLASSIFIED</p> <p>received 670 rad whole body X radiation and orthotopic tail grafts of LAF<sub>1</sub>, BALB/c, C3D/2 and rat skin. Certain groups of mice received specific antisera intraperitoneally, produced in response to BALB/c or rat spleen cells or to two consecutive BALB/c or rat skin grafts. The data indicate that the first-set response to an allogenic skin graft can be significantly inhibited, in sublethally irradiated mice, by specific antisera, while the first-set response to a xenogenic skin graft remains resistant to similar treatment. Specific antisera had no effect upon a pre-existing second-set response. The significance of these data is discussed.</p> <p>UNCLASSIFIED</p>
---	---	--	--

Naval Radiological Defense Laboratory  
USNRDL-TR-625

DIFFERENTIAL EFFECTS OF SPECIFIC ANTISERA  
ON THE REJECTION OF ALLOGENIC AND XENO-  
GENIC SKIN GRAFTS BY SUBLETHALLY X-IRRADI-  
ATED MICE by M. L. Tyan and L. J. Cole  
4 March 1963 19 p. tables 14 refs.

UNCLASSIFIED

Twelve to fourteen week old  
female LAF<sub>1</sub> mice, non-sensitized  
or pre-sensitized with two consecu-  
tive BALB/c or rat skin grafts,  
(over)

1. Immune serums.
2. Skin.
3. Transplantation.
4. Radiation tolerance.
5. Mice.
6. Rats.

- I. Tyan, M. L.
- II. Cole, L. J.
- III. Title.
- IV. NR005.08-5200

UNCLASSIFIED

received 670 rad whole body X radiation and orthotopic tail grafts of LAF<sub>1</sub>. BALB/c, C3D/2 and rat skin. Certain groups of mice received specific antisera intraperitoneally, produced in response to BALB/c or rat spleen cells or to two consecutive BALB/c or rat skin grafts. The data indicate that the first-set response to an allogenic skin graft can be significantly inhibited, in sublethally irradiated mice, by specific antisera, while the first-set response to a xenogenic skin graft remains resistant to similar treatment. Specific antisera had no effect upon a pre-existing second-set response. The significance of these data is discussed.

UNCLASSIFIED